



Structural Racism & Tobacco

BACKGROUND

Structural racism is the normalization and legitimization of historical, cultural, institutional, and interpersonal dynamics that advantageously benefit White people while producing adverse outcomes for people of color.¹ As a result, communities of color are limited in their opportunities for social, economic, and financial advancement ultimately cultivating negative health consequences and inequities, particularly among Black Americans who are also subjected to anti-Black racism.¹ U.S. public policies have systematically displaced and destabilized communities of color, most notably Black and Indigenous communities. Exclusionary policies such as redlining, the Indian Relocation Act of 1956, and the formation of Chinatowns resulted in racially segregated, under resourced, environmentally hazardous, and poverty-concentrated neighborhoods that continue to persist today.^{1,2}

The experience of racism results in chronic discrimination and stress that adversely impacts the health of individuals from historically racially marginalized populations.¹ The use of tobacco products can further exacerbate these negative health consequences resulting in even worse health outcomes among these populations. One study demonstrated that adolescents that had interpersonal experiences with racism were more likely to initiate smoking, regardless of gender, ethnicity, or socioeconomic status.³ Racially or ethnically marginalized communities are less likely to have access to affordable health care, lower medical access and coverage for tobacco cessation, and live in areas with higher tobacco retailer density.⁴⁻⁶ The tobacco industry has an extensive history of targeting historically under-resourced communities with tactics that use racial profiling, neighborhood demographics, and cultural elements to promote tobacco sales.⁷⁻⁹ Additionally, tobacco companies have heavily marketed menthol products to Black communities and low-income neighborhoods.⁸ Strategic targeting by the tobacco industry in combination with structurally racist policies has resulted in disproportionate rates of tobacco use and exposure in historically under-resourced communities.

The American Heart Association is committed to a tobacco and nicotine endgame that ultimately leads to an end to all tobacco and nicotine addiction in the US for all communities. We support first minimizing the use of all combustible tobacco products, while combating the structural barriers that continue to inhibit equitable health outcomes. We aim to ensure the next generation of youth and adolescents do not become addicted to harmful nicotine products.

KEY FACTS

- American Indian/Alaska Native (34.9%) adults had the highest prevalence of tobacco use compared to White (21.1%), Black (19.4%), Hispanic (11.7%), and Asian (11.5%) adults in 2020.¹⁰
- Sovereign tribal status and cultural practices are important factors when considering strategies to address the high prevalence of tobacco use among American Indians/Alaska Natives. Tobacco companies have historically targeted these communities by tactics such as price reductions, giveaways, promotions, charitable contributions, and sponsorships.¹¹
- Black and Hispanic communities have been traditionally pursued by the tobacco industry. These communities experience high levels of tobacco retailer density, price discounts, availability of little cigar and cigarillo, and advertising.^{7,12}
- Menthol cigarettes are used at disproportionately higher rates by racial and ethnic minority smokers, including African Americans (84.6%), Hispanics or Latinos (46.9%) and Asian Americans (38%), compared to White smokers (28.9%).¹³
- In 2020, 37% of teen users of flavored tobacco used menthol flavored products, including 52% of Native American/Native Alaskan flavored tobacco users.¹⁴

FACT SHEET: Structural Racism & Tobacco

- A 2018 study examining the promotion of alternative tobacco products in New York City found that inexpensive, combustible, and most harmful tobacco products are disproportionately more accessible and advertised in non-White and low-income neighborhoods. Black and Hispanic neighborhoods were more likely to carry inexpensive products such as 99-cent cigarillos.¹⁵
- Disproportionately high distribution of tobacco retailers in Black and Hispanic neighborhoods provides increased availability of tobacco products in these communities contributing to existing tobacco disparities. Historical factors such as redlining, racially biased retailer decisions to invest (or not invest) in resources, and neighborhood segregation could all be contributing factors to high retailer density in these communities.¹⁶
- It is important to consider the traditional and cultural difference among population subgroups when targeting different tobacco cessation and control strategies as smoking behavior can vary widely. For example, Asian Americans have the lowest prevalence of cigarette use among racial groups; however, 20% of Koreans smoke cigarettes compared to 7.6% of Chinese individuals.¹⁷
- In the most recent pooled data (2014 – 2017), Native Hawaiian/Other Pacific Islander (23.4%) and American Indians/Alaska Natives (20.6%) youth had the highest prevalence of current tobacco use compared to Multiracial (16.5%), White (15.3%), Hispanic (14.6%), Black (11.5%), and Asian (5.0%) youth.¹⁸
- Between 2014–2017, cigars are more frequently used among Black youth, whereas e-cigarettes are the most common product for all other racial/ethnic youth.¹⁸
- Historically marginalized racial communities are less likely to be screened for tobacco use, receive physician advice to quit smoking, and receive tobacco cessation treatment.^{19,20} These disparities in cessation behaviors could possibly be associated with difference in tobacco use behaviors, health care access, access to cessation treatments, and lack of awareness of these treatments.²¹
- In 2018, NHANES estimated that 48% of Black non-smokers are exposed to secondhand smoke compared to 22% of White non-smokers and 16.6% of Mexican American non-smokers.²² Additionally, 44.7% of those living below the poverty level and 35.6% of those who live in rental housing are exposed to secondhand smoke compared to 21.4% living above the poverty level and 19.2% who own their housing.²²

References

1. Churchwell K, Elkind MSV, Benjamin RM, Carson AP, Chang EK, Lawrence W, Mills A, Odum TM, Rodriguez CJ, Rodriguez F, Sanchez E, Sharrief AZ, Sims M, Williams O, American Heart Association. Call to Action: Structural Racism as a Fundamental Driver of Health Disparities: A Presidential Advisory From the American Heart Association. *Circulation*. 2020;142(24):e454–e468.
2. Zonta M. Racial Disparities in Home Appreciation. *Center for American Progress*. 2019.
3. Read UM, Karamanos A, João Silva M, Molaodi OR, Enayat ZE, Cassidy A, Cruickshank JK, Harding S. The influence of racism on cigarette smoking: Longitudinal study of young people in a British multiethnic cohort. *PLoS One*. 2018;13(1):e0190496.
4. Hostetter M, Klein S. In Focus: Reducing Racial Disparities in Health Care by Confronting Racism. 2018.
5. Cox LS, Okuyemi K, Choi WS, Ahluwalia JS. A review of tobacco use treatments in U.S. ethnic minority populations. *American journal of health promotion: AJHP*. 2011;25(5 Suppl):S11-30.
6. Ribisl KM, D'Angelo H, Feld AL, Schleicher NC, Golden SD, Luke DA, Henriksen L. Disparities in tobacco marketing and product availability at the point of sale: Results of a national study. *Preventive Medicine*. 2017;105:381–388.
7. Cruz TB, Rose SW, Lienemann BA, Byron MJ, Meissner HI, Baezconde-Garbanati L, Huang L-L, Carroll DM, Soto C, Unger JB. Pro-tobacco marketing and anti-tobacco campaigns aimed at vulnerable populations: A review of the literature. *Tobacco Induced Diseases*. 2019;17:68.
8. Mills SD, Henriksen L, Golden SD, Kurtzman R, Kong AY, Queen TL, Ribisl KM. Disparities in retail marketing for menthol cigarettes in the United States, 2015. *Health & Place*. 2018;53:62–70.
9. D'Silva J, O'Gara E, Villaluz NT. Tobacco industry misappropriation of American Indian culture and traditional tobacco. *Tobacco Control*. 2018;27(e1):e57–e64.
10. Cornelius ME. Tobacco Product Use Among Adults — United States, 2020. *MMWR. Morbidity and Mortality Weekly Report*. 2022;71.
11. Lempert LK, Glantz SA. Tobacco Industry Promotional Strategies Targeting American Indians/Alaska Natives and Exploiting Tribal Sovereignty. *Nicotine & Tobacco Research: Official Journal of the Society for Research on Nicotine and Tobacco*. 2019;21(7):940–948.
12. Iglesias-Rios L, Parascandola M. A historical review of R.J. Reynolds' strategies for marketing tobacco to Hispanics in the United States. *American Journal of Public Health*. 2013;103(5):e15-27.
13. Villanti AC, Mowery PD, Delnevo CD, Niaura RS, Abrams DB, Giovino GA. Changes in the prevalence and correlates of menthol cigarette use in the USA, 2004–2014. *Tobacco Control*. 2016;25(Suppl 2):ii14–ii20.
14. Office on Smoking and Health, National Center for Chronic Disease Prevention and Health Promotion. Historical NYTS Data and Documentation | CDC. 2022.
15. Giovenco DP, Spillane TE, Merizier JM. Neighborhood Differences in Alternative Tobacco Product Availability and Advertising in New York City: Implications for Health Disparities. *Nicotine & Tobacco Research: Official Journal of the Society for Research on Nicotine and Tobacco*. 2019;21(7):896–902.

FACT SHEET: Structural Racism & Tobacco

16. Lee JGL, Sun DL, Schleicher NM, Ribisl KM, Luke DA, Henriksen L. Inequalities in tobacco outlet density by race, ethnicity and socioeconomic status, 2012, USA: results from the ASPIRE Study. *Journal of Epidemiology and Community Health*. 2017;71(5):487–492.
17. Martell BN, Garrett BE, Caraballo RS. Disparities in Adult Cigarette Smoking - United States, 2002-2005 and 2010-2013. *MMWR. Morbidity and mortality weekly report*. 2016;65(30):753–758.
18. Odani S. Racial/Ethnic Disparities in Tobacco Product Use Among Middle and High School Students — United States, 2014–2017. *MMWR. Morbidity and Mortality Weekly Report*. 2018;67.
19. Tibuakuu M, Okunrintemi V, Jirru E, Echouffo Tcheugui JB, Orimoloye OA, Mehta PK, DeFilippis AP, Blaha MJ, Michos ED. National Trends in Cessation Counseling, Prescription Medication Use, and Associated Costs Among US Adult Cigarette Smokers. *JAMA network open*. 2019;2(5):e194585.
20. Babb S. Disparities in Cessation Behaviors Between Hispanic and Non-Hispanic White Adult Cigarette Smokers in the United States, 2000–2015. *Preventing Chronic Disease*. 2020;17.
21. Creamer MR, Wang TW, Babb S, Cullen KA, Day H, Willis G, Jamal A, Neff L. Tobacco Product Use and Cessation Indicators Among Adults — United States, 2018. *MMWR. Morbidity and Mortality Weekly Report*. 2019;68(45):1013–1019.
22. Shastri SS, Talluri R, Shete S. Disparities in Secondhand Smoke Exposure in the United States: National Health and Nutrition Examination Survey 2011-2018. *JAMA Internal Medicine*. 2021;181(1):134–137.